## **INFUSION ORDERS: COVID THERAPY: REMDESIVIR**

Name:			DOB:		
Date of symptom onset:_		_ Medication Aller	gies:		
Height:	inches Weight:	kg P	atient Phone #:		
12 and older weighing at	least 40 kg) with <u>pos</u> pitalization or death. I	<u>itive direct SARS-C</u> Please mark the cr	CoV-2 viral testing who a	re at high risk	and pediatric patients (age of progression to severe gh risk category. Failure to
Pregnancy Diabetes Chronic kidney disc	v/growthcharts/clinica ease	al_charts.htm)		er based on C	CDC growth charts,
Immunosuppressiv					
Cardiovascular disease Chronic lung disease Sickle cell disease Neurodevelopment Dependence on me	se (COPD, moderate	to severe asthma,	,		
COVID-19])		-g, (	, gae,, e. pee	о регосовно се	
*** Note patients shoul	d be symptomatic ar	nd within 7 days (	of symptom onset***		
REMDESIVIR TREATME	ENT:				
					lesivir 100 mg in 250 mL s during administration.
X After infusion is com	plete, flush the tubi	ng with sodium c	nloride 0.9% injection t	o ensure del	ivery of the entire dose.
X Observe patient for a	at least 1 hour after i	infusion is compl	ete for infusion-related	reactions.	
X Adverse Reaction/A	naphylaxis Protocol	if necessary (refe	r to form PO500-022-N	i-1).	
X Ondansetron ODT 4	mg sublingual x 1 as	s needed for naus	sea.		
My sig	gnature indicates th	nat I have discus	sed the risks and ber	nefits of this	therapy.
Provider Sign:		_(Print):	J	Date:	Time:
			pleted order to 406-4	147-2719	
DATIENT IDENTIFICA	TIONI			St. Peter's 1	Health
PATIENT IDENTIFICAT	HON.		· ·	30. I CIEI 8 1	iicaitii

**INFUSION ORDERS: COVID THERAPY:** 

2475 Broadway • Helena, MT 59601 (406) 442-2480

PO500-077-N-1 (4-2022)